

Characterization of anisotropic Liouville-type mixed spaces and its application

V. A. Nogin and E. E. Ournycheva

Abstract

Within the framework of the method of hypersingular integrals we obtain a characterization of the anisotropic spaces $L_{\bar{p}, \bar{r}}^{\bar{\alpha}}$. These spaces are defined to consist of functions $f(x) \in L_{\bar{r}}$ for which

$$F^{-1} \left(\sum_{j=1}^n |\xi_j|^{\alpha_j} \right) Ff \in L_{\bar{p}}, \quad \alpha_j > 0.$$

The mentioned characterization is applied to prove the denseness of the class C_0^∞ in $L_{\bar{p}, \bar{r}}^{\bar{\alpha}}$.

2000 Mathematics Subject Classification: Primary 46E35, 47G10; Secondary 42B10, 42B15.

1 Introduction

We consider the anisotropic Liouville-type mixed spaces, which are defined via Fourier transform as follows:

$$L_{\bar{p}, \bar{r}}^{\bar{\alpha}} \equiv L_{\bar{p}, \bar{r}}^{\bar{\alpha}}(\mathbb{R}^n) = \left\{ f : f \in L_{\bar{r}}, F^{-1} \left(\sum_{j=1}^n |\xi_j|^{\alpha_j} \right) Ff \in L_{\bar{p}} \right\}, \quad (1.1)$$

where $\alpha_i > 0$, $\bar{p} = (p_1, \dots, p_n)$, $\bar{r} = (r_1, \dots, r_n)$, $1 \leq p_i, r_i < \infty$ ($i = 1, \dots, n$), $L_{\bar{p}} \equiv L_{\bar{p}}(\mathbb{R}^n)$ being the well-known space equipped with the mixed norm

$$\|f\|_{\bar{p}} = \left(\int_{\mathbb{R}^1} \left(\int_{\mathbb{R}^1} \dots \left(\int_{\mathbb{R}^1} |f(x)|^{p_1} dx_1 \right)^{\frac{p_2}{p_1}} \dots dx_{n-1} \right)^{\frac{p_n}{p_{n-1}}} dx_n \right)^{\frac{1}{p_n}}. \quad (1.2)$$

(We note that some basic properties of the spaces $L_{\bar{p}}$ were established in [1, 2].) The Fourier transform Ff in (1.1) is treated in the sense of distributions on a suitable space of test functions (see Section 2).

In the isotropic case when $\alpha_1 = \dots = \alpha_n = \alpha$, $p_1 = \dots = p_n = p$, and $r_1 = \dots = r_n = r$ the spaces (1.1) coincide with the spaces $L_{p,r}^\alpha$ of functions $f \in L_r$ having their Riesz derivatives $\mathbb{D}^\alpha f \in L_p$. Such spaces were first introduced and investigated by S. G. Samko in [13, 14] (see also the books [16] and [17]). We note that the spaces (1.1) were treated by P. I. Lizorkin in the case $\bar{p} = \bar{r}$ (see [8]–[10]). The spaces $L_{p,r}^{\bar{\alpha}}$ were studied by A. A. Davtayan in [3, 4] in the case $1 < p < n/\alpha^*$, $1/\alpha^* = \frac{1}{n} \sum_{j=1}^n 1/\alpha_j$, $1 < r < \infty$, and by V. A. Nogin and G. P. Emgusheva (see [5, 6]) for $1 \leq p, r < \infty$.

Here we deal with the most general anisotropic case of vector-valued $\bar{\alpha}$, $\bar{p}$, and $\bar{r}$. We obtain a characterization of the spaces $L_{\bar{p},\bar{r}}^{\bar{\alpha}}$ via anisotropic hypersingular integrals (HSI)

$$(T^{\bar{\alpha}}f)(x) = \lim_{\substack{\varepsilon \rightarrow 0 \\ (L_{\bar{p}})}} (T_\varepsilon^{\bar{\alpha}}f)(x), \quad (1.3)$$

introduced by P. I. Lizorkin (see [9]), where $T_\varepsilon^{\bar{\alpha}}f$ is the "truncated" integral

$$(T_\varepsilon^{\bar{\alpha}}f)(x) = \int_{\rho(t) > \varepsilon} \frac{(\Delta_t^{2\ell}f)(x)}{\rho^{n+\alpha^*}(t)} dt, \quad 2\ell > \max_j \alpha_j. \quad (1.4)$$

Here $(\Delta_t^\ell f)(x)$ is the centered finite difference of the function $f(x)$. The function $\rho(t)$, which is referred to as "anisotropic distance", is a positive solution of the equation $\sum_{i=1}^n x_i^2 \rho^{-2\lambda_i} = 1$, $\lambda_i = \alpha^*/\alpha_i$. Namely, we prove that

$$L_{\bar{p},\bar{r}}^{\bar{\alpha}} = \{f : f \in L_{\bar{r}}, \quad T^{\bar{\alpha}}f \in L_{\bar{p}}\}, \quad (1.5)$$

$\alpha_i > 0$, $1 < p_i < \infty$, $1 \leq r_i < \infty$, under some additional restrictions on $\bar{\alpha}$ in the case $\alpha^* \geq n$ (see (2.1)).

We also give an application of the mentioned characterization. Basing ourselves on equality (1.5), we prove that the class C_0^∞ is dense in $L_{\bar{p},\bar{r}}^{\bar{\alpha}}$, $\alpha_i > 0$, $1 < p_i < \infty$, $1 \leq r_i < \infty$ (under the mentioned restrictions on $\bar{\alpha}$ in the case $\alpha^* \geq n$). It should be noted that the denseness of the class C_0^∞ in $L_{p,r}^\alpha$ was proved in [13] for $1 < p < n/\alpha$ and in [11] for $p \geq n/\alpha$. In the

anisotropic case the corresponding result was established in [12], where the authors assumed that $r_i > p_i$ or $r_i \leq p_i$ for every i , $1 \leq i \leq n$, if $\sum_{i=1}^n \frac{1}{\alpha_i p_i} \leq 1$. One of the goals of this paper is to get rid of these unnatural restrictions.

We note that the proof of denseness given in Theorem 2.2 looks very non-trivial. We first prove the statement of this theorem in the case $0 < \max_j \alpha_j < 1$. After that, we extend it to the remaining values of parameters with the aid of induction.

We also point out the papers [19] and [20], where the author dealt with some generalization of the spaces of $L_{p,r}^{\bar{\alpha}}$ -type to the case of a more general anisotropic distance (in comparison with that used in (1.4)).

The paper is organized as follows. In Section 2 we formulate our main results, see Theorems 2.1 and 2.2. Sections 3 and 4 can be regarded as a background to the proof of the mentioned theorems. They contain the necessary preliminaries and some auxiliary statements respectively. In Section 5 we prove Theorems 2.1 and 2.2. For the sake of convenience we gather some technical fragments of the proofs in Appendix. Some results presented in this paper were announced in [12].

2 The main results

We define the Liouville-type spaces $L_{\bar{p},\bar{r}}^{\bar{\alpha}}$, $1 \leq r_i < \infty$, $1 \leq p_i < \infty$, $\alpha_i > 0$, $i = 1, \dots, n$, by equality (1.1), where the Fourier transform Ff is interpreted in the sense of Ψ' -distributions (see Subsection 3.1). We put

$$\|f\|_{L_{\bar{p},\bar{r}}^{\bar{\alpha}}} = \|f\|_{\bar{r}} + \left\| F^{-1} \left(\sum_{j=1}^n |\xi_j|^{\alpha_j} \right) Ff \right\|_{\bar{p}}.$$

The following theorem provides a characterization of the space $L_{\bar{p},\bar{r}}^{\bar{\alpha}}$ by means of HSI (1.3).

Theorem 2.1. *Let $1 < p_i < \infty$, $1 \leq r_i < \infty$, $\alpha_i > 0$, $i = 1, \dots, n$, and*

$$\sum_{j=1}^n \frac{1+k_j}{\alpha_j} \neq 1, \quad |k| = 0, 1, \dots, m-1, \quad m = [\gamma],$$

$$\gamma = \max_j \alpha_j \left(1 - \sum_{j=1}^n \frac{1}{\alpha_j} \right). \quad (2.1)$$

Then equality (1.5) holds. Moreover, the norms $\|f\|_{L_{\vec{p}, \vec{r}}^{\vec{\alpha}}}$ and $\|f\|_{\vec{r}} + \|T^{\vec{\alpha}}f\|_{\vec{p}}$ are equivalent.

Remark 2.1. For the rest of the paper we assume the condition (2.1) to be fulfilled.

We apply the characterization (1.5) to prove the next theorem.

Theorem 2.2. Let $1 < p_i < \infty$, $1 \leq r_i < \infty$, $\alpha_i > 0$, $i = 1, \dots, n$. Then the space C_0^∞ is dense in $L_{\vec{p}, \vec{r}}^{\vec{\alpha}}$.

3 Preliminaries

3.1 Notation:

S is the Schwartz space of rapidly decreasing smooth functions; C_0^∞ is its subspace of finite functions; Ψ_0 , Ψ are the Lizorkin spaces of functions in S , vanishing together with all their derivatives at the origin and on the coordinate hyperplanes respectively; Φ_0 , Φ are their Fourier duals; Ψ' , Φ'_0 are the corresponding spaces of distributions; $C_0^\infty(0, \infty)$ is the class of finite C^∞ -functions on the semi-axis $(0, \infty)$ supported outside zero; $M_{\vec{p}}$ is the class of $\vec{p}$ -multipliers; $X \rightarrow Y$ is the continuous embedding of the normed space X into the normed space Y ; $(\Delta_y^\ell f)(x) = \sum_{k=0}^{\ell} (-1)^k C_\ell^k f(x + (\ell/2 - k)y)$ is the centered finite difference of the function $f(x)$ of order ℓ with the step y ; $(\tilde{\Delta}_y^\ell f)(x) = \sum_{k=0}^{\ell} (-1)^k C_\ell^k f(x - ky)$ is the non-centered finite difference; $(Ff)(y) = \hat{f}(y) = \int_{\mathbb{R}^n} f(x) e^{ix \cdot y} dx$ is the Fourier transform of the function $f(x)$; $(F^{-1}f)(x) = \tilde{f}(x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} f(y) e^{-ix \cdot y} dy$ is the inverse Fourier transform; $\langle f, \omega \rangle = \int_{\mathbb{R}^n} \overline{f(x)} \omega(x) dx$. All constants in various estimates not necessarily the same at each occurrence are denoted by the same letter C . The end of proof is denoted by ■.

3.2 Estimates for some integrals and finite differences

The following lemmas were proved in [13] in the isotropic situation (when $\rho(x) = |x|$, $p_1 = \dots = p_n$, and $s_1 = \dots = s_n$). The proofs of their analogues in the anisotropic case are much in lines with those given in [13] and we omit them.

Lemma 3.1. *Let $\gamma > 0$, $1 \leq s_i < \infty$, $i = 1, \dots, n$,*

$$I(t) = \left(\int_{\mathbb{R}^1} dx_n \dots \left(\int_{\mathbb{R}^1} \left(\prod_{k=0}^{\ell} [1 + \rho(x - kt)]^{-\gamma s_1} \right) dx_1 \right)^{s_2/s_1} \dots \right)^{1/s_n},$$

$\ell = 1, 2, \dots$. Then

$$I(t) \leq C(1 + \rho(t))^{\sum_{i=1}^n \lambda_i/s_i - (\ell+1)\gamma}, \quad \text{if } \frac{1}{\ell+1} \sum_{i=1}^n \frac{\lambda_i}{s_i} < \gamma < \sum_{i=1}^n \frac{\lambda_i}{s_i}, \quad (3.1)$$

$$I(t) \leq C(1 + \rho(t))^{-\gamma\ell}, \quad \text{if } \gamma > \sum_{i=1}^n \frac{\lambda_i}{s_i}. \quad (3.2)$$

Lemma 3.2. *If $a(x) \in C_0^\infty$, then the following inequality holds:*

$$\left| (\tilde{\Delta}_t^\ell a)(x) \right| \leq C \left[\frac{\rho^\ell(t)}{\prod_{k=0}^{\ell} (1 + \rho(x - kt))} \right]^\theta, \quad \ell = 1, 2, \dots, \quad \theta = \min_i \left(\frac{\alpha_i^*}{\alpha_i} \right). \quad (3.3)$$

$x, t \in \mathbb{R}^n$.

Lemma 3.3. *If $a(x) \in C_0^\infty$, then*

$$\left\| \tilde{\Delta}_t^\ell a \right\|_{\bar{p}} \leq C \left[\frac{\rho(t)}{1 + \rho(t)} \right]^{\ell\theta}, \quad \ell = 1, 2, \dots, \quad \theta = \min_i \left(\frac{\alpha_i^*}{\alpha_i} \right). \quad (3.4)$$

3.3 Some properties of the mixed spaces $L_{\bar{p}}$

We need the following properties of functions in $L_{\bar{p}}$.

Theorem 3.1 ([2]). *Let $f(x) \in L_{\bar{p}}(\mathbb{R}^n)$, $1 \leq p_i < \infty$, $i = 1, \dots, n$, and let $a(t)$ be an averaging kernel, that is, $a(t) \in L_1$ and $\int_{\mathbb{R}^n} a(t) dt = 1$. Then*

$$\lim_{\delta \rightarrow 0} \|f - f_\delta\|_{\bar{p}} = 0,$$

where

$$f_\delta(x) = \int_{\mathbb{R}^n} a(t) f(x - \delta t) dt. \quad (3.5)$$

The following anisotropic interpolation inequality “for m points” was proved in [18].

Theorem 3.2. *Let $\bar{p}^j = (p_1^j, \dots, p_n^j)$, $1 \leq p_i^j \leq \infty$, $i = 1, \dots, n$, $j = 1, \dots, m$, $m \leq n + 1$, and $f \in \bigcap_{j=1}^m L_{\bar{p}^j}$, $\theta_k \geq 0$, $\sum_{k=1}^m \theta_k = 1$. Then $f \in L_{\bar{p}}$ for any $\bar{p}$ such that $1/p_i = \sum_{j=1}^m \theta_j/p_i^j$, $i = 1, \dots, n$, and*

$$\|f\|_{\bar{p}} \leq \|f\|_{\bar{p}^1}^{\theta_1} \|f\|_{\bar{p}^2}^{\theta_2} \dots \|f\|_{\bar{p}^m}^{\theta_m}. \quad (3.6)$$

3.4 Anisotropic potentials

Let $\mathcal{K}_{\bar{\alpha}}(x)$ be a smooth function in $\mathbb{R}^n \setminus \{0\}$ such that $\mathcal{K}_{\bar{\alpha}}(t^\lambda x) = t^{\alpha^* - n} \mathcal{K}_{\bar{\alpha}}(x)$, where $x \in \mathbb{R}^n$, $\lambda \in \mathbb{R}_+^n$, and $t > 0$ (that is, the function $\mathcal{K}_{\bar{\alpha}}(x)$ is λ -homogeneous of order $\alpha^* - n$).

An operator of the form

$$(K_{\bar{\alpha}} \varphi)(x) = \int_{\mathbb{R}^n} \mathcal{K}_{\bar{\alpha}}(x - y) \varphi(y) dy \quad (3.7)$$

is called *anisotropic potential* (of order α^*). This operator is well-defined on the whole space $L_{\bar{p}}$, $1 \leq p_i \leq \infty$ ($i = 1, \dots, n$), if $\sum_{i=1}^n \frac{1}{\alpha_i p_i} > 1$. The following theorem can be regarded as an anisotropic analogue of the well-known Sobolev theorem for the Riesz potential operator.

Theorem 3.3 ([10]). *Let $1 < p_i \leq q_i \leq \infty$ ($i = 1, \dots, n-1$), $1 < p_n < q_n < \infty$, $\sum_{i=1}^n \frac{1}{\alpha_i p_i} > 1$, and*

$$\sum_{i=1}^n \frac{1}{\alpha_i q_i} = \sum_{i=1}^n \frac{1}{\alpha_i p_i} - 1. \quad (3.8)$$

Then the operator $K^{\bar{\alpha}}$ is bounded from $L_{\bar{p}}$ into $L_{\bar{q}}$.

Remark 3.1. *The statement of Theorem 3.3 can be generalized to the case of λ -homogeneous kernels of some order $\gamma < 0$ (not necessarily $\gamma = \alpha^* - n$). One can obtain such a generalization replacing (3.8) by the following condition*

$$\sum_{i=1}^n \frac{1}{\alpha_i q_i} = \sum_{i=1}^n \frac{1}{\alpha_i p_i} - \frac{\gamma + n}{\alpha^*}.$$

For $\sum_{i=1}^n \frac{1}{\alpha_i p_i} \leq 1$ we treat the potential $K^{\bar{\alpha}}\varphi$, $\varphi \in L_{\bar{p}}$, in the sense of Φ'_0 -distributions:

$$\langle K^{\bar{\alpha}}\varphi, \omega \rangle = \langle \varphi, K^{\bar{\alpha}}\omega \rangle, \quad \omega \in \Phi_0, \quad \varphi \in L_{\bar{p}}. \quad (3.9)$$

Such a treatment is correct since $\mathcal{K}_{\bar{\alpha}}(x)$ is a convolute in Φ_0 , in accordance with the Gel'fand-Shilov theorem (see [7], P.155), because $\widehat{\mathcal{K}_{\bar{\alpha}}}(\xi)$ is a multiplier in Ψ_0 .

We consider some anisotropic potentials of special-type. Let $Q^{\bar{\alpha}}$ be the operator with the kernel $Q_{\bar{\alpha}}(x) = F^{-1} \left(\frac{1}{S_{\bar{\alpha}}(\xi)} \right) (x)$, where $S_{\bar{\alpha}}(\xi)$ is a symbol of HSI (1.3). The kernel $Q_{\bar{\alpha}}(x)$ admits the following representations (see [4] and [6] in the cases $\alpha^* < n$ and $\alpha^* \geq n$ respectively):

$$Q_{\bar{\alpha}}(x) = \int_0^\infty t^{n-\alpha^*-1} \tilde{g}(t^\lambda x) dt \quad , \text{ if } \quad \alpha^* < n, \quad (3.10)$$

$$\begin{aligned} Q_{\bar{\alpha}}(x) &= m \sum_{k=|m|} \frac{x^k}{k!} \int_0^1 t^{n-\alpha^*+\lambda \cdot k-1} dt \int_0^1 (1-u)^{m-1} (D^k \tilde{g})(ut^\lambda x) du + \\ &+ \int_1^\infty t^{n-\alpha^*-1} \tilde{g}(t^\lambda x) dt, \quad m = [\gamma] + 1, \\ \gamma &= (\alpha^* - n) \max_j \frac{\alpha_j}{\alpha^*} \quad , \text{ if } \quad \alpha^* \geq n, \end{aligned}$$

where $g(\xi) = \varphi(\rho(\xi))/S_{\bar{\alpha}}(\xi)$, $\varphi \in C_0^\infty(0, \infty)$, $\varphi(t) \geq 0$, and $\int_0^\infty \frac{\varphi(t)}{t} dt = 1$.

The function $Q_{\bar{\alpha}}(x)$ is λ -homogeneous of order $\alpha^* - n$, if $\alpha^* < n$. In the case $\alpha^* \geq n$ it is λ -homogeneous in the Φ'_0 -sense, that is,

$$t^{\alpha^*-n} \langle Q_{\bar{\alpha}}(x/t^\lambda), \omega \rangle = \langle Q_{\bar{\alpha}}, \omega(t^\lambda x) \rangle, \quad \omega \in \Phi_0.$$

We also consider the modified potential $\mathcal{R}^{\bar{\alpha}}\varphi$ with the kernel $\mathcal{R}_{\bar{\alpha}}(x)$, being λ -homogeneous (in the regular sense) for any $\alpha^* > 0$, which differs from $Q_{\bar{\alpha}}(x)$ by a polynomial in the case $\alpha^* \geq n$ (see [6]):

$$\begin{aligned} \mathcal{R}_{\bar{\alpha}}(x) &= Q_{\bar{\alpha}}(x), & \text{if } \alpha^* < n, \\ \mathcal{R}_{\bar{\alpha}}(x) &= Q_{\bar{\alpha}}(x) + \sum_{|k| \leq m-1} \frac{x^k}{k!} \frac{(D^k \tilde{g})(0)}{\lambda \cdot k + n - \alpha^*}, & \text{if } \alpha^* \geq n. \end{aligned} \quad (3.11)$$

We observe that

$$\mathcal{R}^{\bar{\alpha}}\varphi = Q^{\bar{\alpha}}\varphi, \quad \varphi \in L_{\bar{p}}, \quad (3.12)$$

in the case $\alpha^* \geq n$, when both potentials are treated in the Φ'_0 -sense, in view of the fact that Φ_0 -functions are orthogonal to all polynomials.

4 Some auxiliary statements

4.1 Characterization of the space $L_{\bar{p}, \bar{r}}^{\bar{\alpha}}$ in terms of anysotropic potentials

Let $Q^{\bar{\alpha}}(L_{\bar{p}})$ be the space of anysotropic potentials:

$$Q^{\bar{\alpha}}(L_{\bar{p}}) = \{f : f = Q^{\bar{\alpha}}\varphi, \varphi \in L_{\bar{p}}\},$$

$$\bar{\alpha} = (\alpha_1, \dots, \alpha_n), \quad \bar{p} = (p_1, \dots, p_n), \quad \alpha_i > 0, \quad 1 \leq p_i < \infty;$$

we set

$$\|f\|_{Q^{\bar{\alpha}}(L_{\bar{p}})} = \|\varphi\|_{\bar{p}}.$$

Theorem 4.1. *Let $1 < p_i < \infty$, $1 \leq r_i < \infty$, $\alpha_i > 0$, $i = 1, \dots, n$. Then*

$$L_{\bar{p}, \bar{r}}^{\bar{\alpha}} = L_{\bar{r}} \cap Q^{\bar{\alpha}}(L_{\bar{p}}).$$

Moreover, the norms $\|f\|_{L_{\bar{p}, \bar{r}}^{\bar{\alpha}}}$ and $\|f\|_{\bar{r}} + \|f\|_{Q^{\bar{\alpha}}(L_{\bar{p}})}$ are equivalent.

Proof. Let $f \in L_{\bar{p}, \bar{r}}^{\bar{\alpha}}$. We observe that

$$f = Q^{\bar{\alpha}}B\varphi, \tag{4.1}$$

where $\varphi = F^{-1} \left(\sum_{j=1}^n |\xi_j|^{\alpha_j} \right) Ff \in L_{\bar{p}}$, B is the bounded operator in $L_{\bar{p}}$, generated by the $\bar{p}$ -multiplier $b(\xi) = S_{\bar{\alpha}}(\xi) / \sum_{j=1}^n |\xi_j|^{\alpha_j}$ (the relation $b(\xi) \in M_{\bar{p}}$ is verified with the aid of Lizorkin theorem, see [10]). Indeed, for $\omega \in \Phi$ we have

$$\langle f, \omega \rangle = \frac{1}{(2\pi)^n} \langle \hat{f}, \hat{\omega} \rangle = \frac{1}{(2\pi)^n} \langle \hat{\varphi}, b(\xi) \frac{\hat{\omega}(\xi)}{S_{\bar{\alpha}}(\xi)} \rangle = \langle B\varphi, Q^{\bar{\alpha}}\omega \rangle.$$

From here we derive equality (4.1), which yields $f \in Q^{\bar{\alpha}}(L_{\bar{p}})$. Relatively, $L_{\bar{p}, \bar{r}}^{\bar{\alpha}} \subset L_{\bar{r}} \cap Q^{\bar{\alpha}}(L_{\bar{p}})$ and $\|f\|_{\bar{r}} + \|f\|_{Q^{\bar{\alpha}}(L_{\bar{p}})} \leq C\|f\|_{L_{\bar{p}, \bar{r}}^{\bar{\alpha}}}$.

Let now $f \in L_{\bar{r}}$ and $f = Q^{\bar{\alpha}}\varphi$, $\varphi \in L_{\bar{p}}$. Then $F^{-1} \left(\sum_{j=1}^n |\xi_j|^{\alpha_j} \right) Ff = B^{-1}\varphi \in L_{\bar{p}}$. Therefore $f \in L_{\bar{p}, \bar{r}}^{\bar{\alpha}}$ and $\|f\|_{L_{\bar{p}, \bar{r}}^{\bar{\alpha}}} \leq C(\|f\|_{\bar{r}} + \|f\|_{Q^{\bar{\alpha}}(L_{\bar{p}})})$. ■

Corollary 4.1. *The spaces $L_{\bar{p}, \bar{r}}^{\bar{\alpha}}$ are complete.*

4.2 On a continuous imbedding of the spaces $L_{\bar{p}, \bar{r}}^{\bar{\alpha}}$ with respect to $\bar{\alpha}$

Theorem 4.2. *Let $1 \leq r_i < \infty$, $1 < p_i < \infty$, $\alpha_i > 0$, and $\beta_i > 0$ be such that $0 < \alpha^* - \beta^* < n$ and $\sum_{i=1}^n \frac{1}{\alpha_i p_i} > 1 - \frac{\beta^*}{\alpha^*}$. Then*

$$L_{\bar{p}, \bar{r}}^{\bar{\alpha}} \longrightarrow L_{\bar{s}, \bar{r}}^{\bar{\beta}}, \quad (4.2)$$

$\bar{s} = (s_1, \dots, s_n)$ being an arbitrary vector such that $s_i \geq p_i$, if $i = 1, \dots, n-1$, $s_n > p_n$, and

$$\sum_{i=1}^n \frac{1}{\alpha_i s_i} = \sum_{i=1}^n \frac{1}{\alpha_i p_i} - \left(1 - \frac{\beta^*}{\alpha^*}\right).$$

Proof. Let us consider the anisotropic potential $I^{\bar{\alpha}}\varphi = k_{\bar{\alpha}} * \varphi$ with kernel $k_{\bar{\alpha}}(x) \stackrel{(\Phi'_0)}{=} F^{-1}(\rho^{-\alpha^*}(\xi))(x)$, which is possessed of the same properties, as the kernel $Q_{\bar{\alpha}}(x)$ (see (3.10)).

Since both functions $\rho^{\alpha^*}(\xi)/S_{\bar{\alpha}}(\xi)$ and $\rho^{-\alpha^*}(\xi)S_{\bar{\alpha}}(\xi)$ belong to $M_{\bar{p}}$ by the Lizorkin theorem (see [10]), we have

$$Q^{\bar{\alpha}}(L_{\bar{p}}) = I^{\bar{\alpha}}(L_{\bar{p}}) \stackrel{\text{def}}{=} \{f : f = I^{\bar{\alpha}}\varphi, \varphi \in L_{\bar{p}}\}.$$

Therefore

$$L_{\bar{p}, \bar{r}}^{\bar{\alpha}} = L_{\bar{r}} \cap I^{\bar{\alpha}}(L_{\bar{p}}) \quad (4.3)$$

by Theorem 4.1.

Let $f = I^{\bar{\alpha}}\varphi$, $\varphi \in L_{\bar{p}}$, and $\omega \in \Phi_0$. We have

$$\langle f, \omega \rangle = \frac{1}{(2\pi)^n} \langle \hat{\varphi}, \rho^{-\alpha^*}(\xi) \hat{\omega}(\xi) \rangle = \langle \psi, I^{\beta} \omega \rangle,$$

where $\psi = F^{-1}(\rho^{\beta^* - \alpha^*}(\xi)) * \varphi$. Making use of Remark 3.1, we easily obtain the imbedding $Q^{\bar{\alpha}}(L_{\bar{p}}) \rightarrow I^{\bar{\beta}}(L_{\bar{s}})$, which yields (4.2) by (4.3). ■

5 Proof of the main results

5.1 Proof of Theorem 2.1

Let $f \in L_{\bar{p}, \bar{p}}^{\bar{\alpha}}$, then $f = Q^{\bar{\alpha}}\varphi$, $\varphi \in L_{\bar{p}}$, by Theorem 4.1. Relatively, $f = \mathcal{R}^{\bar{\alpha}}\varphi$.

We prove the following integral representation

$$(T_{\varepsilon}^{\bar{\alpha}}\mathcal{R}^{\bar{\alpha}}\varphi)(x) = \int_{\mathbb{R}^n} \mathcal{K}(y)\varphi(x - \varepsilon^{\lambda}y)dy, \quad (5.1)$$

where the kernel

$$\mathcal{K}(y) = \rho^{-n}(y) \int_{\rho(t) > 1/\rho(y)} \frac{(\Delta_t^{2\ell}\mathcal{R}_{\bar{\alpha}})\left(\frac{y}{\rho^{\lambda}(y)}\right)}{\rho^{n+\alpha^*}(t)} dt \quad (5.2)$$

is averaging. We note that the representation (5.1) was proved in [6] for $\varphi \in \Phi_0$.

Since the symbol

$$S_{\bar{\alpha}, \varepsilon}(\xi) = (-1)^{\ell} 2^{2\ell} \int_{\rho(t) > \varepsilon} \frac{\sin^{2\ell} \frac{\xi \cdot t}{2}}{\rho^{n+\alpha^*}(t)} dt$$

of the "truncated" integral (1.4) is a multiplier in Ψ_0 (see [9]), we conclude that the space Φ_0 is invariant with respect to the operator $T_{\varepsilon}^{\bar{\alpha}}$. Therefore

$$\begin{aligned} \langle T_{\varepsilon}^{\bar{\alpha}}f, \omega \rangle &= \langle f, T_{\varepsilon}^{\bar{\alpha}}\omega \rangle = \langle \varphi, \mathcal{R}^{\bar{\alpha}}T_{\varepsilon}^{\bar{\alpha}}\omega \rangle = \langle \varphi, T_{\varepsilon}^{\bar{\alpha}}\mathcal{R}^{\bar{\alpha}}\omega \rangle = \langle \varphi, \int_{\mathbb{R}^n} \mathcal{K}(y)\omega(x - \varepsilon^{\lambda}y)dy \rangle = \\ &= \langle \int_{\mathbb{R}^n} \mathcal{K}(y)\varphi(x - \varepsilon^{\lambda}y)dy, \omega \rangle, \quad \omega \in \Phi_0. \end{aligned}$$

Thus (5.1) is valid in the Φ'_0 -sense.

Since two locally integrable functions, which agree in the Φ'_0 -sense, may differ from each other only by a polynomial, we obtain (5.1) for almost all $x \in \mathbb{R}^n$ (because both sides of (5.1) belong to the weighted spaces $L_{1, \gamma} = \{f(x) : \int_{\mathbb{R}^n} |f(x)| (1 + |x|)^{\gamma} dx < \infty\}$ for some $\gamma > -n$).

Letting $\varepsilon \rightarrow 0$ in (5.1), in view of Theorem 3.1 we have

$$T^{\bar{\alpha}}f = \varphi \in L_{\bar{p}}, \quad \|T^{\bar{\alpha}}f\|_{\bar{p}} = \|f\|_{Q^{\bar{\alpha}}(L_{\bar{p}})}.$$

Let now $f \in L_{\bar{r}}$, $T^{\bar{\alpha}}f \in L_{\bar{p}}$. The statement of theorem will follow from the relation $f = Q^{\bar{\alpha}}\varphi$, where $\varphi = T^{\bar{\alpha}}f$. For $\omega \in \Phi_0$ we have

$$\begin{aligned} \langle \varphi, Q^{\bar{\alpha}}\omega \rangle &= \langle \lim_{\substack{\varepsilon \rightarrow 0 \\ (L_{\bar{p}})}} T_{\varepsilon}^{\bar{\alpha}}f, \mathcal{R}^{\bar{\alpha}}\omega \rangle = \lim_{\varepsilon \rightarrow 0} \langle T_{\varepsilon}^{\bar{\alpha}}f, \mathcal{R}^{\bar{\alpha}}\omega \rangle = \\ &= \lim_{\varepsilon \rightarrow 0} \langle f, \int_{\mathbb{R}^n} \mathcal{K}(y)\omega(x - \varepsilon^\lambda y)dy \rangle = \langle f, \omega \rangle. \end{aligned} \quad (5.3)$$

The second of equalities of this chain follows from the fact that the convergence in $L_{\bar{p}}$ implies that in Φ'_0 , the last one is justified by the Hölder inequality. Thus we have proved that

$$\langle f, \omega \rangle = \langle \varphi, Q^{\bar{\alpha}}\omega \rangle. \quad (5.4)$$

In the case $\sum_{i=1}^n \frac{1}{\alpha_i p_i} > 1$ we have

$$\langle f, \omega \rangle = \langle Q^{\bar{\alpha}}\varphi, \omega \rangle$$

hence, $f = Q^{\bar{\alpha}}\varphi$ for almost all $x \in \mathbb{R}^n$. In the case $\sum_{i=1}^n \frac{1}{\alpha_i p_i} \leq 1$ the relation (5.4) itself means that $f = Q^{\bar{\alpha}}\varphi$. Now the application of Theorem 4.1 yields $f \in L_{\bar{p}, \bar{r}}^{\bar{\alpha}}$. ■

Corollary 5.1. *Let $f \in L_{\bar{p}, \bar{r}}^{\bar{\alpha}}$, $\alpha_i > 0$, $1 < p_i < \infty$, $1 \leq r_i < \infty$, $i = 1, \dots, n$. Then the following integral representation holds:*

$$(\Delta_t^m f)(x) = \int_{\mathbb{R}^n} (\Delta_t^m \mathcal{R}_{\bar{\alpha}})(x - y)(T^{\bar{\alpha}}f)(y)dy, \quad m > \max_j \alpha_j. \quad (5.5)$$

Corollary 5.2. *Let $f \in L_{\bar{p}, \bar{r}}^{\bar{\alpha}}$, $\alpha_i > 0$, $1 < p_i < \infty$, $1 \leq r_i < \infty$ ($i = 1, \dots, n$). Then*

$$\|\Delta_t^m f\|_{\bar{p}} \leq c\rho^{\alpha^*}(t)\|T^{\bar{\alpha}}f\|_{\bar{p}}, \quad m > \max_j \alpha_j. \quad (5.6)$$

We give the proofs of Corollaries 5.1 and 5.2 in Appendix.

5.2 Proof of Theorem 2.2

We first approximate the function $f \in L_{\overline{p}, \overline{r}}^\alpha$ by smooth functions using the averages (3.5) with a "very nice" kernel. Namely, we assume that $a(t) \in C_0^\infty$. Then $\|f - f_\delta\|_{L_{\overline{p}, \overline{r}}^\alpha} \rightarrow 0$, as $\delta \rightarrow 0$. Therefore it remains to approximate $f \in C^\infty \cap L_{\overline{p}, \overline{r}}^\alpha$ by C_0^∞ -functions.

For this goal, we consider the function $\mu(x) \in C_0^\infty$ supported in the ball $\rho(x) < 2$ such that $0 \leq \mu(x) \leq 1$, $\mu(x) = 1$ if $\rho(x) < 1$, and $\mu(x) = 0$ in the case $\rho(x) \geq 2$. Let $\mu_N(x) = \mu(N^{-\lambda}x)$. We show that the smooth truncations $f_N(x) = \mu_N(x)f(x) (\in C_0^\infty)$ approximate $f(x)$ in the norm of $L_{\overline{p}, \overline{r}}^\alpha$, as $N \rightarrow \infty$. Let $\nu(x) = 1 - \mu(x)$, $\nu_N(x) = \nu(x/N)$. We have to prove that

$$\|T^{\overline{\alpha}}(\nu_N f)\|_{\overline{p}} \rightarrow 0, \quad \text{as } N \rightarrow \infty, \quad f \in C^\infty \cap L_{\overline{p}, \overline{r}}^\alpha, \quad (5.7)$$

where

$$(T^{\overline{\alpha}} f)(x) = \int_{\mathbb{R}^n} \frac{(\Delta_t^{2\ell} f)(x)}{\rho^{n+\alpha^*}(t)} dt, \quad 2\ell > \max_j \alpha_j. \quad (5.8)$$

It should be noted that the integral (5.8) converges absolutely for the functions $f(x)$ bounded in $\mathbb{R}^n$ together with all their partial derivatives up to the order $[\alpha^*] + 1$.

I. Proof of (5.7) in the case $\sum_{i=1}^n \frac{1}{\alpha_i p_i} > 1$.

We first suppose that $0 < \max_j \alpha_j < 1$ and take $\ell = 1$ in (5.8). Then

$$|T^{\overline{\alpha}}(\nu_N f)(x)| \leq |\nu_N(x) (T^{\overline{\alpha}} f)(x)| + 2 |(B_N f)(x)|, \quad (5.9)$$

where

$$(B_N f)(x) = \int_{\mathbb{R}^n} \frac{\nu_N(x-t) - \nu_N(x)}{\rho^{n+\alpha^*}(t)} f(x-t) dt.$$

In order to obtain (5.7), it suffices to verify that

$$\|B_N f\|_{\overline{p}} \rightarrow 0, \quad N \rightarrow \infty \quad (5.10)$$

(since $\|\nu_N(x)f(x)\|_{\overline{p}} \rightarrow 0$, as $N \rightarrow \infty$, for every $f(x) \in L_{\overline{p}}$).

We justify (5.10) for $f \in L_{\bar{q}}$, where $\bar{q}$ is an arbitrary vector such that $\bar{q} > \bar{p}$ and equality (3.8) is fulfilled. This implies (5.10) in view of the imbedding $L_{\bar{p}, \bar{r}}^{\bar{\alpha}} \subset L_{\bar{q}}$, which is valid by virtue of Theorems 4.1 and 3.3.

Owing to the uniform estimate

$$\|B_N f\|_{\bar{p}} \leq C \|f\|_{\bar{q}}, \quad (5.11)$$

with the constant C not depending on N (see Appendix), in accordance with the Banach–Steinhaus theorem, it remains to verify (5.10) for $f \in C_0^\infty$ (since the class C_0^∞ is dense in $L_{\bar{q}}$).

Assuming that $f(x) \in C_0^\infty$ is supported in the ball $\rho(x) < a$, we have

$$\begin{aligned} \|B_N f\|_{\bar{p}} &\leq \frac{C}{N^\theta} + C N^{\alpha^* \sum_{i=1}^n \frac{1}{\alpha_i p_i} - \alpha^*} \int_{\rho(t) > \frac{2a}{C_1 N}} \frac{dt}{\rho^{n+\alpha^*-\theta}(t)} \times \\ &\times \left(\int_{\mathbb{R}^1} dx_n \dots \left(\int_{\mathbb{R}^1} \frac{|f(N^\lambda x)|^{p_1} dx_1}{(1+\rho(x))^{p_1 \theta} (1+\rho(x-t))^{p_1 \theta}} \right)^{\frac{p_2}{p_1}} \dots \right)^{\frac{1}{p_n}}, \end{aligned}$$

where $\theta = \alpha^* / \max_j \alpha_j$, C_1 being such that $\rho(x-t) \geq C_1 \rho(t) - \rho(x)$. Since $\rho(x-t) \geq C_1 \rho(t) - \frac{a}{N} > 0$, if $\rho(x) < \frac{a}{N}$ and $\rho(t) > \frac{2a}{C_1 N}$, we have

$$\|B_N f\|_{\bar{p}} \leq \frac{C}{N^\theta} + \frac{C}{N^{\alpha^*}} \int_{\mathbb{R}^n} \frac{dt}{\rho^{n+\alpha^*-\theta}(t) \left(1 + C_1 \rho(t) - \frac{a}{N_0}\right)^\theta} \rightarrow 0,$$

as $N \rightarrow \infty$ ($N > N_0 > a$). Thus we have obtained the desired result in the case $0 < \max_j \alpha_j < 1$.

In order to prove (5.7) in the case $\max_j \alpha_j \geq 1$, we use the induction argument. It should be noted that such an idea is due to S.G. Samko (see [13]) in the isotropic case. Its realization in the most general anisotropic case of vector-valued $\bar{\alpha}$, $\bar{p}$, and $\bar{r}$, treated here, is much more difficult. The following lemma, which enables one us to apply the induction argument, plays a crucial role in the proof of Theorem 2.2.

Lemma 5.1. *Let $m = 1, 2, \dots$ and $\sum_{i=1}^n \frac{1}{\alpha_i p_i} > 1$. If the class C_0^∞ is dense in $L_{\bar{p}, \bar{r}}^{\bar{\alpha}}$ for $0 < \max_j \alpha_j < m$, then it is also dense in $L_{\bar{p}, \bar{r}}^{\bar{\alpha}}$ for $m \leq \max_j \alpha_j < m+1$.*

Proof. As above, we have to check (5.7) for $m \leq \max_j \alpha_j < m + 1$.

Owing to the evident formulas $(\Delta_t^{2\ell} \nu_N f)(x) = (\tilde{\Delta}_t^{2\ell} \nu_N f)(x + \ell t)$,
 $(\tilde{\Delta}_t^{2\ell} \nu_N f)(x) = \sum_{k=0}^{2\ell} C_{2\ell}^k (\tilde{\Delta}_t^k \nu_N)(x) (\tilde{\Delta}_t^{2\ell-k} f)(x - kt)$, we have

$$T^{\bar{\alpha}}(\nu_N f) = B_N f + \sum_{k=1}^{2\ell} C_{2\ell}^k B_{N,k} f,$$

where

$$(B_N f)(x) = \int_{\mathbb{R}^n} \frac{\nu_N(x + \ell t) (\tilde{\Delta}_t^{2\ell} f)(x + \ell t)}{\rho^{n+\alpha^*}(t)} dt;$$

$$(B_{N,k} f)(x) = \int_{\mathbb{R}^n} \frac{(\tilde{\Delta}_t^k \nu_N)(x + \ell t) (\tilde{\Delta}_t^{2\ell-k} f)(x + (\ell - k)t)}{\rho^{n+\alpha^*}(t)} dt,$$

$$k = 1, 2, \dots, 2\ell.$$

Let us show that

$$\|B_{N,k} f\|_{\bar{p}} \rightarrow 0, \quad \text{as } N \rightarrow \infty, \quad k = 1, 2, \dots, 2\ell, \quad (5.12)$$

and

$$\|B_N f\|_{\bar{p}} \rightarrow 0, \quad \text{as } N \rightarrow \infty. \quad (5.13)$$

In the case $k = 2\ell$ the relation (5.12) is proved for $f \in L_{\bar{q}} (\supset L_{\bar{p}, \bar{r}}^{\bar{\alpha}})$ in just the same way, as (5.7) for $0 < \max_j \alpha_j < 1$.

In the cases $k = 1, 2, \dots, 2\ell - 1$ we prove the stronger relation:

$$\|B_{N,k} f\|_{\bar{p}} \rightarrow 0, \quad \text{as } N \rightarrow \infty, \quad f \in L_{\bar{\tau}^k, \bar{r}}^{\bar{\gamma}^k}, \quad (5.14)$$

the vectors $\bar{\gamma}^k = (\gamma_1^k, \dots, \gamma_n^k)$ and $\bar{\tau}^k = (\tau_1^k, \dots, \tau_n^k)$ being such that $\sum_{i=1}^n \frac{1}{\alpha_i \tau_i^k} =$

$$\sum_{i=1}^n \frac{1}{\alpha_i p_i} - \left(1 - \frac{(\gamma^k)^*}{\alpha^*}\right), \quad \tau_i^k > p_i, \quad \text{and}$$

$$0 < \max_i \gamma_i^k < m. \quad (5.15)$$

(Later we will put some additional restrictions on γ_i^k). Then (5.14) yields (5.12) by (4.2) for $f \in L_{\bar{p}, \bar{r}}^\alpha$ and $\max_i \gamma_i^k < m \leq \max_i \alpha_i$. We first obtain the uniform estimate

$$\|B_{N,k}f\|_{\bar{p}} \leq C\|f\|_{L_{\bar{\tau}^k, \bar{r}}^{\bar{\gamma}^k}}, \quad f \in L_{\bar{\tau}^k, \bar{r}}^{\bar{\gamma}^k}. \quad (5.16)$$

Making use of (3.3) and the Hölder inequality, we obtain

$$\begin{aligned} \|B_{N,k}f\|_{\bar{p}} &\leq \frac{C}{N^{k\theta}} \int_{\mathbb{R}^n} \frac{\|\tilde{\Delta}_t^{2\ell-k}f\|_{\bar{\tau}^k}}{\rho^{n+\alpha^*-k\theta}(t)} dt \times \\ &\quad \times \left(\int_{\mathbb{R}^1} dx_n \dots \left(\int_{\mathbb{R}^1} \frac{dx_1}{\prod_{i=0}^k \left(1 + \frac{\rho(x+(\ell-i)t)}{N}\right)^{\theta\xi_1^k}} \right)^{\xi_2^k/\xi_1^k} \dots \right)^{1/\xi_n^k}, \end{aligned}$$

where $\xi_i^k = \frac{p_i\tau_i^k}{\tau_i^k - p_i}$. Applying Lemma 3.2, under the additional condition $2\ell - k > \max_j \gamma_j^k$, we have

$$\|\Delta_t^{2\ell-k}f\|_{\bar{\tau}^k} \leq c\rho^{(\gamma^k)^*}(t) \|T^{\bar{\gamma}^k}f\|_{\bar{\tau}^k}, \quad f \in L_{\bar{\tau}^k, \bar{r}}^{\bar{\gamma}^k},$$

where

$$\sum_{i=1}^n \frac{1+j_i}{\gamma_i^k} \neq 1, \quad |j| = 0, 1, \dots, \left[\max_i \gamma_i^k \left(1 - \sum_{i=1}^n \frac{1}{\gamma_i^k}\right) \right] - 1. \quad (5.17)$$

Besides this, we assume that

$$\gamma_i^k < \frac{(2\ell - k)\alpha_i}{\max_j \alpha_j}, \quad i = 1, \dots, n, \quad (5.18)$$

then

$$\begin{aligned} \|B_{N,k}f\|_{\bar{p}} &\leq C \|T^{\bar{\gamma}^k}f\|_{\bar{\tau}^k} \int_{\mathbb{R}^n} \frac{dt}{\rho^{n+\alpha^*-k\theta-(\gamma^k)^*}} \times \\ &\quad \times \left(\int_{\mathbb{R}^1} dx_n \dots \left(\int_{\mathbb{R}^1} \frac{dx_1}{\prod_{i=0}^k [1 + \rho(x + (l-i)t)]^{\theta\xi_1^k}} \right)^{\xi_2^k/\xi_1^k} \dots \right)^{1/\xi_n^k}. \end{aligned}$$

In order to apply Lemma 3.1, we also put the following restrictions on $\bar{\gamma}^k$:

$$(\gamma^k)^* > \alpha^* - \theta k. \quad (5.19)$$

Then Lemma 3.1 yields

$$\|B_{N,k}f\|_{\bar{p}} \leq C \left\| T^{\bar{\gamma}^k} f \right\|_{\bar{\tau}^k}. \quad (5.20)$$

Taking into account (5.15), (5.18), and (5.19), we arrive at the following inequalities for the coordinates γ_i^k :

$$\max \left(0, \alpha_i - \frac{\alpha_i k}{\max_j \alpha_j} \right) < \gamma_i^k < \min \left(\frac{m \alpha_i}{\max_j \alpha_j}, \frac{(2\ell - k) \alpha_i}{\max_j \alpha_j} \right). \quad (5.21)$$

Since $\max_j \alpha_j < 2\ell$ and $m \leq \max_j \alpha_j < m + 1$, this interval is not empty.

Thus we have proved (5.16). In accordance with the Banach–Steinhaus theorem, it suffices to verify (5.14) on a dense set in $L_{\bar{\tau}^k, \bar{r}}^{\bar{\gamma}^k}$. Since $0 < \max_i \gamma_i^k < m$, the class C_0^∞ is dense in $L_{\bar{\tau}^k, \bar{r}}^{\bar{\gamma}^k}$ according to the assumption of induction. For $f(x) \in C_0^\infty$, in view of (3.3) we have

$$\begin{aligned} \|B_{N,k}f\|_{\bar{p}} &\leq \frac{C}{N^{(\gamma^k)^* - (2\ell - k)\theta}} \int_{\mathbb{R}^n} \frac{dt}{\rho^{n+\alpha^* - 2\ell\theta}(t)(1 + N\rho(t))^{(2\ell - k)\theta}} \times \\ &\times \left(\int_{\mathbb{R}^1} dx_n \dots \left(\int_{\mathbb{R}^1} \frac{dx_1}{\prod_{i=0}^k (1 + \rho(x - it))^{\theta \xi_1^k}} \right)^{\xi_2^k / \xi_1^k} \dots \right)^{1/\xi_n^k}. \end{aligned}$$

Applying the Hölder inequality to the inner integral, we get

$$\|B_{N,k}f\|_{\bar{p}} \leq \frac{c}{N^{(\gamma^k)^* - (2\ell - k)\theta + d}} \int_{\mathbb{R}^n} \frac{dt}{\rho^{n+\alpha^* - 2\ell\theta + d}(t)(1 + \rho(t))^{(k+1)\theta - \alpha^* + (\gamma^k)^*}}, \quad (5.22)$$

where $0 \leq d \leq (2\ell - k)\theta$. The integral on the right-hand side converges, if we take $d \in ((2\ell - k)\theta - (\gamma^k)^*, \min\{(2\ell - k)\theta, 2\ell\theta - \alpha^*\})$ (this interval is not empty in view of (5.19)). Then the right-hand side of (5.22) tends to zero, as $N \rightarrow \infty$.

Let us prove (5.13). We have

$$(B_N f)(x) = \nu_N(x) (T^{\bar{\alpha}} f)(x) + (M_N f)(x),$$

where

$$(M_N f)(x) = \int_{\mathbb{R}^n} \frac{\nu_N(x + \ell t) - \nu_N(x)}{\rho^{n+\alpha^*}(t)} \left(\tilde{\Delta}_t^{2\ell} f \right)(x + \ell t) dt. \quad (5.23)$$

The validity of (5.13) follows from the relation

$$\|M_N f\|_{\bar{p}} \rightarrow 0, \quad \text{as } N \rightarrow \infty, \quad f \in L_{\bar{p}, \bar{r}}^{\bar{\alpha}},$$

which is verified in just the same way, as (5.12).

II. The case $\sum_{i=1}^n \frac{1}{\alpha_i p_i} \leq 1$.

In the case $0 < \max_j \alpha_j < 1$, as above, we have (5.9).

Let $p_j < r_j$ for $j = m_1^1, \dots, m_{k_1}^1$ and $p_j \geq r_j$ for $j = m_1^2, \dots, m_{k_2}^2$, and let $\bar{r}^1$ be the vector with the following coordinates: $r_j^1 = r_j$ for $j = m_1^1, \dots, m_{k_1}^1$ and $r_j^1 = p_j$ for $j = m_1^2, \dots, m_{k_2}^2$. We observe that $f \in L_{\bar{r}^1}$ by virtue of (3.6). Similarly to (5.10) we prove that

$$\|B_N f\|_{\bar{p}} \rightarrow 0, \quad \text{as } N \rightarrow \infty, \quad f \in L_{\bar{r}^1}. \quad (5.24)$$

Making use of the Minkowsky and Hölder inequalities, we arrive at the estimate

$$\|B_N f\|_{\bar{p}} \leq N^{-\alpha^* + \alpha^* \sum_{j=m_1^1}^{m_{k_1}^1} \frac{1}{\alpha_j p_j} - \alpha^* \sum_{j=m_1^1}^{m_{k_1}^1} \frac{1}{\alpha_j r_j}} \|f\|_{\bar{r}^1} \int_{\mathbb{R}^n} \frac{\|\nu(x - t) - \nu(x)\|_{\bar{s}}}{\rho^{n+\alpha^*}(t)} dt,$$

where $s_j = \frac{p_j r_j}{r_j - p_j}$ for $j = m_1^1, \dots, m_{k_1}^1$ and $s_j = \infty$ for $j = m_1^2, \dots, m_{k_2}^2$.

Since $\alpha^* - \alpha^* \sum_{j=m_1^1}^{m_{k_1}^1} \frac{1}{\alpha_j p_j} + \alpha^* \sum_{j=m_1^1}^{m_{k_1}^1} \frac{1}{\alpha_j r_j} > 0$ and the integral on the right-hand side converges, we obtain (5.24).

In order to extend the result on denseness of C_0^∞ in $L_{\bar{p}, \bar{r}}^{\bar{\alpha}}$ to the case $\max_j \alpha_j \geq 1$, we use the following analogue of Lemma 5.1.

Lemma 5.2. *Let $m = 1, 2, \dots$, $1 < p_i < \infty$, $1 \leq r_i < \infty$, $\alpha_i > 0$, and $\sum_{i=1}^n \frac{1}{\alpha_i p_i} \leq 1$. If C_0^∞ is dense in $L_{\bar{p}, \bar{r}}^{\bar{\alpha}}$ for $0 < \max_j \alpha_j < m$, then it is dense in $L_{\bar{p}, \bar{r}}^{\bar{\alpha}}$ for $m \leq \max_j \alpha_j < m + 1$.*

The proof of this lemma is much in lines with that of Lemma 5.1. The only difference is as follows. To prove (5.12), we have to use the assumption of induction only in the case $\frac{1}{\max_j \alpha_j} - \sum_{j=m_1^1}^{m_{k_1}^1} \frac{1}{\alpha_j p_j} + \sum_{j=m_1^1}^{m_{k_1}^1} \frac{1}{\alpha_j r_j} \leq 0$ and $k = 1, \dots, [\max_j \alpha_j]$. As in the proof of Lemma 5.1, we base ourselves on the imbedding (4.2). Namely, we choose β_i ($i = 1, \dots, n$) such that $\alpha^* - \beta^* < \min\{n, \theta\}$, $0 < \max_j \beta_j < m$, $\sum_{i=1}^n \frac{1 + j_i}{\beta_i} \neq 1$,

$|j| = 0, 1, \dots, \left[\max_i \beta_i \left(1 - \sum_{i=1}^n \frac{1}{\beta_i} \right) \right] - 1$, and check (5.12) for $f \in L_{\bar{\zeta}, \bar{r}}^{\bar{\beta}}$, where the components of $\bar{\zeta}$ ($\zeta_i > p_i$, $i = 1, \dots, n$) satisfy the equality $\sum_{i=1}^n \frac{1}{\zeta_i \alpha_i} = \sum_{i=1}^n \frac{1}{\alpha_i p_i} - \left(1 - \frac{\beta^*}{\alpha^*} \right)$.

As regards the rest values of k , the relation (5.12) is proved by the direct estimation of the norm $\|B_{N,K} f\|_{\bar{p}}$. We leave the corresponding details to the reader. ■

6 Appendix

6.1 Proof of equality (5.5).

Let $f \in L_{\bar{p}, \bar{r}}^{\bar{\alpha}}$. We denote $\varphi_\varepsilon = T_\varepsilon^{\bar{\alpha}} f$, then

$$\int_{\mathbb{R}^n} (\Delta_t^m \mathcal{R}_{\bar{\alpha}})(x - y) \varphi_\varepsilon(y) dy = \int_{\mathbb{R}^n} (\Delta_t^m f)(x - \varepsilon^\lambda y) \mathcal{K}(y) dy, \quad (6.1)$$

where $\mathcal{K}(y)$ is the kernel (5.2). Let us pass to the limit, as $\varepsilon \rightarrow 0$, in (6.1).

Making use of the estimate

$$(\Delta_{\eta_t}^m \mathcal{R}_{\bar{\alpha}})(y) \leq c \rho^{\alpha^* - n - m \frac{\alpha^*}{\max_j \alpha_j}}(y), \quad \rho(y) > A, \quad \eta_t = t / \rho^\lambda(t)$$

(see [4]), we arrive at the relation $(\Delta_t^m \mathcal{R}_{\bar{\alpha}})(y) \in L_1$. Therefore the left-hand side of (6.1) converges in $L_{\bar{p}}$, as $\varepsilon \rightarrow 0$, to $T^{\bar{\alpha}} f$, while the right-hand side

converges in $L_{\bar{r}}$ to $\Delta_t^m f$, since $f(x) \in L_{\bar{r}}$ and $\mathcal{K}(y) \in L_1$. This implies (5.5). $\blacksquare$

6.2 Proof of inequality (5.6).

With the aid of (5.5) we have

$$\|\Delta_t^m f\|_{\bar{p}} \leq \rho^{\alpha^*}(t) \|T^{\bar{\alpha}} f\|_{\bar{p}} \int_{\mathbb{R}^n} |(\Delta_{\eta_t}^m \mathcal{R}_{\bar{\alpha}})(y)| dy,$$

where $\eta_t = t/\rho^\lambda(t)$. Since the integral on the right-hand side is finite, we obtain (5.6). $\blacksquare$

6.3 Proof of (5.11)

The application of (3.3) yields

$$|(B_N f)(x)| \leq \frac{C}{N^\theta} \int_{\mathbb{R}^n} \frac{\rho^\theta(t) |f(x-t)| dt}{\left(1 + \frac{\rho(x)}{N}\right)^\theta \left(1 + \frac{\rho(x-t)}{N}\right)^\theta \rho^{n+\alpha^*}(t)},$$

where $\theta = \frac{\alpha^*}{\max_j \alpha_j}$. Applying the Minkowsky and Hölder inequalities, we obtain

$$\begin{aligned} \|B_N f\|_{\bar{p}} &\leq C \|f\|_{\bar{q}} \int_{\mathbb{R}^n} \frac{dt}{\rho^{n+\alpha^*-\theta}(t)} \times \\ &\times \left(\int_{\mathbb{R}^1} \dots \left(\int_{\mathbb{R}^1} \left(\int_{\mathbb{R}^1} \frac{dx_1}{(1+\rho(x))^{\theta\beta_1} (1+\rho(x-t))^{\theta\beta_1}} \right)^{\frac{\beta_2}{\beta_1}} dx_2 \right)^{\frac{\beta_3}{\beta_2}} \dots dx_n \right)^{\frac{1}{\beta_n}}, \end{aligned}$$

where $\beta_i = \frac{p_i q_i}{q_i - p_i}$, $i = 1, \dots, n$. Since $\sum_{i=1}^n \frac{1}{\alpha_i \beta_i} < \frac{\theta}{\alpha^*}$, we arrive at (5.11) by virtue of (3.2). $\blacksquare$

Acknowledgments. We are thankful to Prof. S. G. Samko for the useful discussions of the results and to Prof. E. Liflyand for many helpful comments to the manuscript of the paper.

References

- [1] A. BENEDEK, R. PANZONE, The spaces L_p with mixed norm. Duke Math. J., 28, No 3 (1961), 302–324.
- [2] O. V. BESOV, V. P. IL'IN, S. M. NIKOL'SKII, Integral representations of functions and imbedding theorems (Russian). Nauka (Moscow, 1975). Transl. Vol. 1, John & Sons, New-York (1978).
- [3] A. A. DAVTYAN, Anisotropic potentials, their inversion and some applications (Russian). Dokl. Akad. Nauk SSSR, 285, No 3 (1985), 537–541. Transl. Soviet Math., 32 (1985), 717–721.
- [4] A. A. DAVTYAN, Spaces of anisotropic potentials. Applications (Russian). Trudy Mat. Inst. Steklov, 173 (1986), 113–124. Transl. Proc. Steklov Inst. of Math. 173 (1987), 119–131.
- [5] G. P. EMGUSHEVA, V. A. NOGIN, On the denseness of infinitely differentiable finite functions in the anisotropic classes of Liouville type (Russian). Deposited in VINITI (08.08.88), No 6313–88.
- [6] G. P. EMGUSHEVA, V. A. NOGIN, Characterization of functions in the anisotropic classes of Liouville type (Russian). Izv. Vyssh. Uchebn. Zaved. Math., No 7 (1989), 63–66.
- [7] I. M. GEL'FAND, G. E. SHILOV, Generalized functions, Vol. 2: Fundamental and generalized functions. New York: Acad. Press (1968).
- [8] P. I. LIZORKIN, Generalized Liouville differentiation and the method of multipliers in the theory of imbeddings of classes of differentiable functions (Russian). Trudy Mat. Inst. Steklov, 105 (1969), 89–167. Transl. Proc. Steklov Inst. Math., 105 (1969), 105–202.
- [9] P. I. LIZORKIN, Description of the spaces $L_p^r(\mathbb{R}^n)$ in terms of singular difference integrals (Russian). Matem. sbornik (N. S.), 81(123), No 1 (1970), 79–91. Transl. Math. USSR Sb., 10 (1970), 77–89.

- [10] P. I. LIZORKIN, Multipliers of Fourier integrals and estimates of convolution in spaces with mixed norm. Applications (Russian). *Izv. Akad. Nauk SSSR, Ser. Mat.*, 34 No 1 (1970), 218–247. Transl. Math. USSR *Izv.*, 4 (1970), 225–255.
- [11] V. A. NOGIN, S. G. SAMKO, On the simultaneous approximation of functions and their Riesz derivatives (Russian). *Dokl. Akad. Nauk SSSR*, 261, No 3 (1981), 548–550. Transl. Soviet Math., 24 (1981), 552–554.
- [12] V. A. NOGIN, E. E. OURNYCHEVA, On the denseness of infinitely differentiable finite functions in the anisotropic classes of Liouville type with mixed norms. Collection of papers. *Izd. DGTU (Rostov-on-Don*, 1998), 81–91.
- [13] S. G. SAMKO, On spaces of Riesz potentials (Russian). *Izv. Akad. Nauk SSSR, Ser. Mat.*, 40, No 5 (1976), 1143–1172. Transl. Math. USSR *Izv.*, 10, No 5 (1976), 1089–1117.
- [14] S. G. SAMKO, The spaces $L_{p,r}^\alpha(\mathbb{R}^n)$ and hypersingular integrals (Russian). *Studia Math. (PRL)*, 61, No 3, (1977), 193–230.
- [15] S. G. SAMKO, Denseness of the space Φ_V of Lizorkin type in the mixed $L^{\vec{p}}(\mathbb{R}^n)$ -spaces. *Studia Math. (PRL)*, 113, No 3, (1991), 199–210.
- [16] S. G. SAMKO, A. A. KILBAS, O. I. MARICHEV, Fractional Integrals and Derivatives. Gordon and Breach Science Publishers. (1993).
- [17] S. G. SAMKO, Hypersingular integrals and their applications. Taylor & Francis, Series "Analytical Methods and Special Functions", Vol. 5 (2002).
- [18] S. G. SAMKO, E. E. OURNYCHEVA, The spaces $L_{p_2}^{rad}(L_{p_1}^{ang})$ with radial-spherical mixed norms. *Izv. Vyssh. Uchebn. Zaved., Mat.*, No 4 (1996), 50–59.

- [19] T. A. HAMIDOVA, Hypersingular integrals and spaces of anysotropic Riesz potentials (Russian). Deposited in VINITI (27.12.88), No 9067–88.
- [20] T. A. HAMIDOVA, On denseness of infinitely differentiable finite functions in the anysotropic spaces $Z_{p,r}^\alpha$ (Russian). Deposited in VINITI (28.11.90), No 5976–90.

e-mail: vnogin@math.rsu.ru

e-mail: ournyce@macs.biu.ac.il